

ORLICAINTM

Pat. Pend.



Active Ingredients:

Tetracaine HCL;
Bupivacaine HCL;
Phenylephrine HCL

Purpose:

Oral Anesthetic

Recommended dose:

0.3 milligrams per
kilogram based on FDA
approved drug

Warnings:

For external use
only. DO NOT
INJECT.

Flavors:

Mint, Bubblegum

Uses:

Orlicaine is a flavored topical anesthetic gel providing rapid and profound anesthesia of the oral tissues including the hard and soft palate. The active ingredients fit well within the safety profile of an approved FDA drug and have been studied in children down to the age of three years old.

Directions:

Orlicaine Jar

Orlicaine Mint 14ml, Orlicaine Bubble Gum 14ml

Remove lid. Express directly onto cotton swab applicator or glove. Dry tissue slightly before application. Place Orlicaine on tissue and rinse gently after 2-4 minutes depending on the depth of anesthesia desired.

Anesthesia will occur within 30 seconds and continue to become more profound for up to 15 minutes. Duration ranges from 30 to 90 minutes.

Orlicaine Clinical Kit for SRP

Orlicaine Mint 25ml in a bottle with a luer lock adapter

Remove the black lid and replace with the white luer lock adapter cap. Connect a 1ml luer lock syringe to the adapter cap. Turn the bottle upside down and draw up only the amount of product needed for the procedure. Place the bent, blunt luer lock tip provided on the syringe.

Dry tissue slightly before application. Express Orlicaine directly on the patient's tissue or draw desired amount from luer lock dispenser for syringe delivery into the gingival sulcus using the provided blunt SRP tip. *In the maxillary arch, Orlicaine in the gingival sulcus will provide the most profound effect and deepest anesthesia for SRPs or any other uncomfortable procedures.*

Rinse gently after 2-4 minutes depending on the depth of anesthesia desired. Anesthesia will occur within 30 seconds and continue to become more profound for up to 15 minutes. Duration ranges from 30 to 90 minutes.

The Bennett Pharma Dental Research Team has performed the following with Orlicaine and without an injection:

- Scaling and Root Planing
- Tissue Biopsy
- Suture Placement and Removal
- Implant Uncovering Procedures
- Frenectomy
- and more...

Questions? Let us help.

Dr. Michael Smilanich, DDS

Medical Director, Bennett Pharma

Phone: 952-381-9027

Email: drmike@bennettpharma.com

Email: info@bennett-pharma.com

